

VOLUME, DIAMETER AND THE MINIMAL MASS OF A STATIONARY 1-CYCLE

A. NABUTOVSKY AND R. ROTMAN

ABSTRACT. In this paper we present upper bounds on the minimal mass of a non-trivial stationary 1-cycle. The results that we obtain are valid for all closed Riemannian manifolds. The first result is that the minimal mass of a stationary 1-cycle on a closed n -dimensional Riemannian manifold M^n is bounded from above by $\frac{(n+2)!d}{3}$, where d is the diameter of a manifold M^n . The second result is that the minimal mass of a stationary 1-cycle on a closed Riemannian manifold M^n is bounded from above by $2(n+2)!FillRad(M^n) \leq 2(n+2)!(n+1)n^n\sqrt{n!}(vol(M^n))^{1/n}$, where $FillRad(M^n)$ is the filling radius of a manifold, and $vol(M^n)$ is its volume.

1. Introduction.

In 1983 M. Gromov asked whether there exists a constant $c(n)$ such that the length of a shortest closed geodesic $l(M^n) \leq c(n)(vol(M^n))^{\frac{1}{n}}$, (see [G], p. 135). This problem also appeared as Problem 87 in a list of open problems in Differential Geometry composed by S.-T. Yau ([Y], p. 689, or [SY], p. 297). In the same spirit it might be interesting to know whether there exists $\tilde{c}(n)$ such that $l(M^n) \leq \tilde{c}(n)d$. In this paper we prove the existence of a stationary 1-cycle such that its mass satisfies these inequalities. In fact, our proofs demonstrate the existence of a stationary 1-cycle of a special type which we will call *optimal* such that its mass satisfies these inequalities. Optimal 1-cycles are defined as follows: An optimal 1-cycle is a 1-cycle representable by a subset of M^n that consists of finitely many points $p_1, \dots, p_l$ and a finite collection of geodesic segments between some of these points. Some of these segments can be (possibly trivial) geodesic loops. We also allow several different geodesic segments between two points, as well as several copies of the same geodesic segment (in other words, every segment can have a positive integer multiplicity). Each point p_i should be adjacent to at least one of these segments. For any p_i the number of the geodesic segments adjacent to p_i

counted with their multiplicities is an even positive number. (Here every geodesic loop that starts and ends at p_i is counted twice.) This condition is necessary and sufficient in order for the subset to represent a 1-cycle. Equivalently, we can define an optimal 1-cycle as a 1-cycle represented by a map of a finite multigraph with vertices of positive even degree into M^n such that each edge is mapped into a (possibly non-minimizing) geodesic connecting the images of the adjacent vertices.

An optimal 1-cycle is *stationary* if and only if the following property holds: For any $i = 1, \dots, l$ the sum of unit vectors in $T_{p_i}M^n$ tangent to all geodesic segments meeting at p_i (counted with their multiplicities) is equal to zero. Here each tangent vector is directed from p_i . (The direction from p_i is formally defined as follows. Parametrize the geodesic segment. The direction from p_i coincides with the direction of the first derivative, if the geodesic segment exits p_i , and is the opposite direction, if the geodesic segment enters p_i . It is easy to see that this definition does not depend on the parametrization.) The mass of an optimal 1-cycle is equal to the sum of lengths of images of all its edges. We also obtain an explicit upper bound for the total number of all segments (counted with their multiplicities) in an optimal stationary 1-cycle of mass not exceeding $\tilde{c}(n)\text{diam}(M^n)$ (or $c(n)\text{vol}(M^n)^{\frac{1}{n}}$) in terms of n .

Of course, our estimates would give the estimates on the length of a shortest closed geodesic if the stationary 1-cycle we obtain is realized by a closed geodesic. And, in fact, when M^n is diffeomorphic to the 2-dimensional sphere, this technique produces a closed geodesic, as was observed by Pitts, Calabi and Cao, (see [ClCo]). This fact enabled us to obtain estimates for the length of a shortest closed geodesic on a manifold diffeomorphic to S^2 , (see [NR1]) improving previously known results by C.B. Croke and M. Maeda, (see [C], [Ma]). (Similar results were independently obtained by S. Sabourau in [S1]. Sabourau had also found curvature free upper bounds on the length of a shortest geodesic loop on a compact Riemannian manifold, (see [S2]). Note that a geodesic loop is a stationary 1-cycle if and only if it is a closed geodesic. Also note that the general upper bounds for the length of shortest closed geodesics can be found in [NR]; see also the earlier paper [R]. Those estimates, however, use information about sectional curvature.)

Our techniques were inspired by the geometric measure theory approach to the existence of minimal submanifolds developed by Almgren and Pitts, (see [P]). We

also use an appropriate generalization of obstruction to an extension technique used by Gromov in [G] (see section 1.2 of [G] as well as the proof of Proposition on p. 136 of [G]), and in the case of the Theorem 2 below we use Gromov's upper bound for the filling radius in terms of volume.

One of our starting points is the Almgren and Pitts minimax argument in the geometric measure theory that produces stationary varifolds by considering minima over non-trivial homotopy classes of maps f of S^k into the spaces of cycles on a compact Riemannian manifold of $\max_{t \in S^k} \text{mass}(f(t))$ (see Theorem 4.10 in [P]). (One gets the stationary varifold for a $t \in S^k$ where the maximum of the mass of $g(t)$ is attained for a map g minimizing this maximum over the homotopy class.) Roughly speaking one can get an upper bound for the volume of stationary varifolds obtained by this method as follows: One can start from an arbitrary map f_0 in a non-trivial homotopy class. Denote by M_0 the maximal mass of cycles $f_0(t)$ over $t \in S^k$. Then M_0 will provide an upper bound for the volume of the varifold.

Non-trivial elements of homotopy groups of the space of cycles correspond to non-trivial homology classes of M^n by virtue of the classical result of F. Almgren ([A]) that establishes the isomorphism between the k -th homotopy group of the space of l -dimensional cycles $Z_l(M^n, G)$ in M^n and the $(k+l)$ th homology group of M^n with coefficients in G (for all k, l and $G = \mathbb{Z}$ or $G = \mathbb{Z}_p$ for some p). The underlying geometric idea behind the above correspondence is slicing. For example, consider a smooth function $f : M^n \rightarrow [0, 1]$ such that $f^{-1}(\{0, 1\})$ consists of finitely many points and such that $f^{-1}(t)$ is an $(n-1)$ -dimensional cycle for any t . Then we obtain a slicing of M^n into $(n-1)$ -dimensional cycles $f^{-1}(\{t\}), t \in [0, 1]$ or, in other words, a map from $[0, 1]$ to the space of $(n-1)$ -dimensional cycles in M^n . Since 0 and 1 are mapped to the zero cycle we obtain, in fact, a map of $S^1 = [0, 1]/\{0, 1\}$ into the space $Z_{n-1}(M^n)$. It gives rise to an element of $\pi_1(Z_{n-1}(M^n))$ that corresponds to the fundamental homology class of M^n under the Almgren isomorphism, and therefore is non-trivial. In fact, Almgren started from assigning a $(k+l)$ -dimensional chain (corr. a $(k+l-1)$ -dimensional cycle) in M^n with coefficients on G to a map of $[0, 1]^k$ (or $\partial[0, 1]^k$) into the space of cycles $Z_l(M^n, G)$. A simple modification of this assignment which we will for brevity call *the Almgren correspondence* will be used in this paper, and will be explained in section 4 (in our context).

In this paper we are interested only in one-dimensional cycles. Therefore we are spared a significant amount of technical difficulties arising in geometric measure theory. In particular, we can afford the luxury of considering only 1-cycles made of finitely many Lipschitz curves on M^n (as in [ClCo]). Therefore all of the above ideas from the geometric measure theory become much more obvious and geometric in our situation.

Now, let us denote the minimal mass (=length) of a non-trivial optimal stationary 1-cycle in M^n by $\alpha(M^n)$. Theorems 1 and 2 below establish upper bounds for $\alpha(M^n)$ in terms of the volume $\text{vol}(M^n)$ and the diameter d of the manifold M^n .

Theorem 1. *Let M^n be a closed simply-connected Riemannian manifold of dimension n . Let $q(\leq n)$ denote the minimal dimension i such that $\pi_i(M^n) \neq 0$.*

Then there exists a non-trivial optimal stationary 1-cycle on M^n that consists of at most $\frac{(q+2)!}{2}$ geodesic segments and loops such that its mass does not exceed $\frac{(q+2)!}{3}d$. If $q = 2$ then there exists a stationary 1-cycle of length $\leq 4d$ that is either a closed geodesic or consists of two geodesic loops emanating from the same point p . (Observe, that in the last case the angles made by the tangent vectors of these loops at p are equal, and the bisectors of these angles in $T_p M^n$ are oppositely directed rays.)

Remark. It is well-known (and easy to prove) that if M^n is not simply-connected then there is a closed geodesic on M^n of length $\leq 2d$.

In order to state the next theorem we will need the following definitions and the following result of Gromov (see [G]).

Definition 1. *Let M^n be a manifold topologically imbedded into an arbitrary metric space X . Then its filling radius, denoted $\text{FillRad}(M \subset X)$, is the infimum of $\epsilon > 0$, such that M^n bounds in the ϵ -neighborhood $N_\epsilon(M^n)$, i.e. homomorphism $H_n(M^n, \mathbb{Z}_2) \longrightarrow H_n(N_\epsilon(M^n), \mathbb{Z}_2)$ induced by the inclusion map vanishes.*

Definition 2. *Let M^n be an abstract manifold. Then its filling radius, denoted $\text{FillRad} M^n$ will be $\text{FillRad}(M \subset X)$, where $X = L^\infty(M)$, i.e. the Banach space of bounded Borel functions f on M^n and the imbedding of M^n into X is the map that to each point p of M^n assigns the distance function $p \longrightarrow f_p = d(p, q)$*

Theorem A ([G]). *Let M^n be a closed connected Riemannian manifold. Then $\text{FillRad}M^n \leq (n+1)n^n n!^{\frac{1}{2}} (\text{vol}M^n)^{\frac{1}{n}}$.*

Theorem 2. *Let M^n be a closed Riemannian manifold. Then there exists a non-trivial optimal stationary 1-cycle in M^n made of at most $(n+2)!/2$ geodesic segments and loops such that its mass is bounded from above by $(n+2)!\text{FillRad}M^n \leq (n+2)!(n+1)n!^{\frac{1}{2}} n^n (\text{vol}(M^n))^{\frac{1}{n}}$.*

2. Basic definitions.

First, let us briefly recall some basic notions of (the one-dimensional) geometric measure theory. Our exposition was influenced by the treatment of this subject in [ClCo], [BZ], [P] and [M]. We refer the reader to [M] or [P] for more details.

We say that a 1-dimensional Borel subset A of M^n is *rectifiable* if there exists a Borel subset $S \subset \mathbb{R}^1$ of finite measure and a 1-1 Lipschitz map $f : S \rightarrow M^n$ such that $f(S) \subset A$ and the 1-dimensional Hausdorff measure of $A \setminus f(S)$ is zero. Any such set A can be regarded as a linear functional on the space of C^∞ -smooth 1-forms on M^n : $T_A(\phi) = \int_A \phi$. When A is regarded as the linear functional T_A on 1-forms, it is called a *rectifiable current*. In general, 1-dimensional rectifiable currents are defined as continuous linear functionals on the space of 1-forms on M^n (with the mass topology explained below) that can be represented as countable linear combinations $\sum_{i=1}^\infty a_i T_{A_i}$ with integer coefficients a_i of functionals T_{A_i} corresponding to 1-dimensional rectifiable sets A_i . It is not difficult to see that an arbitrary rectifiable 1-current can be represented as $\sum_{i=1}^\infty a_i T_{M_i}$, where a_i are integer coefficients, and M_i are 1-dimensional submanifolds of M^n , such that $\sum_{i=1}^\infty |a_i| l(M_i) < \infty$, where l denotes the length.

The space of rectifiable 1-currents can be endowed with the weak topology: $T_i \rightarrow T$ if and only if $T_i(\phi) \rightarrow T(\phi)$ for any 1-form ϕ . This space can also be endowed with the mass norm as follows: Define the *mass norm* of a 1-form ω as $\sup_{x \in M^n} \sup_{v \in T_x M^n; \|v\|=1} |\omega(x)(v)|$. The *mass* $M(T)$ of a 1-dimensional rectifiable current T is defined as the supremum over the set of all 1-forms ω on M^n of mass one of $|T(\omega)|$. Note that if A is a C^1 -smooth curve in M^n with a countable set of self-intersections then $M(T_A) = l(A)$. However, if A backtracks over itself then the lengths of its pieces where the backtracking takes place cancel each other, and the mass is strictly less than the length. The mass is a lower-semicontinuous (but not

continuous) functional on the space of rectifiable currents with the weak topology. The topology induced by the mass norm on the space of 1-currents is called the *mass topology*. Observe that a C^1 -smooth path in the space of smooth closed curves in M^n regarded as currents is always continuous in the weak topology but usually NOT continuous in the mass topology. Indeed, the mass of the difference of two C^1 -close but not intersecting C^1 -smooth closed curves is equal to the *sum* of their lengths. (More generally, if A_i are Lipschitz curves in M^n such that the sets of their pairwise intersections and self-intersections have measure zero, then $M(\sum_i a_i T_{A_i}) = \sum_i |a_i| l(A_i)$.) A sequence of 1-currents corresponding to Lipschitz curves γ_i converges to the 1-current corresponding to a Lipschitz curve γ in the mass topology if and only if the mass of the symmetric difference $(\gamma \setminus \gamma_i) \cup (\gamma_i \setminus \gamma)$ converges to zero.

Let T be a 1-dimensional rectifiable current. If for any smooth function f on M^n $T(df) = 0$, then we say that T is a *rectifiable 1-cycle* on M^n . Define $\bar{Z}_1(M^n, Z)$ as the subspace of the space of all 1-dimensional rectifiable currents endowed with the weak topology that consists of all 1-cycles. Consider the space of 1-cycles made of finitely many curves $Z_1(M^n, Z) \subset \bar{Z}_1(M^n, Z)$ that consists of all 1-cycles made of finitely many Lipschitz curves (in other words, we are considering only those cycles that can be represented as $\sum_{i=1}^k a_i T_{A_i}$, where $A_1, \dots, A_k$ are images of $[0, 1]$ under Lipschitz maps to M^n , and $a_i \in Z$.) Any such cycle is an integer linear combination of currents corresponding to *closed* Lipschitz curves in M^n with integer multiplicities. (Note that the above notation is not standard: the notation $Z_1(M^n, Z)$ is usually used for the space that we denote $\bar{Z}_1(M^n, Z)$ in this paper.) The approximation theorem from the geometric measure theory (cf. [M]) implies that $\bar{Z}_1(M^n, Z)$ is the closure of $Z_1(M^n, Z)$ in either weak or mass topology.

Now we would like to introduce some spaces of “nice” 1-cycles that are especially useful for our purposes. Following [ClCo] it is convenient to consider spaces of *parametrized* 1-cycles made of at most k closed curves: Define Γ_k as the space of all k -tuples $(\gamma_1, \dots, \gamma_k)$ of Lipschitz maps of $[0, 1]$ to M^n such that $\sum_{i=1}^k \gamma_i(0) = \sum_{i=1}^k \gamma_i(1)$. Endow Γ_k with the following metric topology: First, because of the Nash embedding theorem we can assume without any loss of generality that M^n is isometrically embedded into the Euclidean space R^N

of a large dimension. Now define the distance $d((\alpha_1, \dots, \alpha_k), (\gamma_1, \dots, \gamma_k))$ as $\max_{i,t} d_{M^n}(\alpha_i(t), \gamma_i(t)) + \sum_{i=1}^k \sqrt{\int_0^1 |\alpha'_i(t) - \gamma'_i(t)|^2 dt}$. It is easy to see that the length functional $l((\gamma_1, \dots, \gamma_k)) = \sum_{i=1}^k l(\gamma_i)$ is a continuous functional on this space. Also, the map $I : \Gamma_k \longrightarrow Z_1(M^n, Z)$ (with the weak topology) given by the formula $I((\gamma_1, \dots, \gamma_k)) = \sum_{i=1}^k T_{\gamma_i}$ is continuous. Denote the image of Γ_k under I by $Z_{(k)}$. We will call $Z_{(k)}$ the space of 1-cycles on M^n made of at most k closed curves, and we will call k the order of these 1-cycles. Observe that Γ_k contains all collections of at most k suitably parametrized closed curves in M^n . Therefore $Z_{(1)} \subset Z_{(2)} \subset Z_{(3)} \subset \dots$, and $\bigcup_{i=1}^{\infty} Z_{(i)} = Z_1(M^n, Z)$. Also, for any x let the subset of Γ_k formed by all $\gamma = (\gamma_1, \dots, \gamma_k)$ such that $l(\gamma) = \sum_{i=1}^k l(\gamma_i) \leq x$ be denoted by Γ_k^x , and the image of Γ_k^x under I be denoted by $Z_{(k)}^x$. We will call $Z_{(k)}^x$ the space of 1-cycles on M^n of length $\leq x$ made of at most k curves. Similarly, we will call elements of Γ_k parametrized 1-cycles made of k curves, k will be called the order of parametrized cycles from Γ_k , and elements of Γ_k^x will be called parametrized 1-cycles of length $\leq x$ made of k curves.

Our definition of *stationary* parametrized or non-parametrized 1-cycles made of k segments is quite similar to the standard definition in the geometric measure theory (cf. [P]): Let X be a vector field on M^n . It determines the one-parameter group of diffeomorphisms $\Phi_X(t)$ of M^n . For any $\gamma \in \Gamma_k$ $\Phi_X(t)(\gamma)$ is a continuous one-parameter family of parametrized 1-cycles. Now we can consider the continuous function $L_{X,\gamma}(t)$ defined as the total length of k Lipschitz curves that together form $\Phi_X(t)(\gamma)$. If for any X $t = 0$ is the critical point of $L_{X,\gamma}$ then γ is called a *stationary parametrized 1-cycle made of k geodesic segments*. Its image $I(\gamma)$ in $Z_{(k)}$ is called a *stationary (non-parametrized) 1-cycle made of k geodesic segments*. The value of the derivative of $L_{X,\gamma}$ at $t = 0$ will be called *the first variation of the length of γ in the direction of X* . In order to understand the stationarity condition we first observe that each of k curves γ_i comprising γ must be a geodesic in a neighborhood of every point $\gamma_i(t)$, where γ_i does not intersect itself or γ_j for some $j \neq i$, if γ is a stationary parametrized 1-cycle. (Otherwise we can use a local vector field X supported on a small ball centered at this point of γ_i to demonstrate that γ is not stationary.) Now we can consider vector fields supported in a small neighborhood of $a = \gamma_i(1)$ or $a = \gamma_i(0)$. When the neighborhood is small the first variation of length is dominated by $\langle X(a), \sum_{i \in I_1} \gamma'_i(1) - \sum_{i \in I_2} \gamma'_i(0) \rangle$, where I_1

is the set of $i \in \{1, \dots, k\}$ such that $\gamma_i(1) = a$, and $I_2 \in \{1, \dots, k\}$ is the set of i such that $\gamma_i(0) = a$. Hence for any such a the sum of the tangent vectors to curves γ_i meeting at a (and directed to a) must be equal to zero. A similar condition must hold at any point $\gamma_i(t)$, where γ_i has an intersection of finite multiplicity with itself and/or other curves γ_j , $j \neq i$. However, in general, γ_i need not be a geodesic in a neighborhood of such a point even if γ is stationary. For example, let $k = 1$. Assume that $\gamma = \gamma_1$ is a three-petal curve that consists of three geodesic loops emanating from the same point p . Consider three angles formed by tangent vectors at p to these three loops. (As usual in this paper, we direct these tangent vectors from p .) Assume further that: 1) These three angles have equal values that are strictly less than $\pi/6$; 2) The bisectors of these three angles lie in a plane in $T_p M^n$ and form angles equal to $\pi/3$ with each other. Then it is easy to see that $\gamma \in \Gamma_1$ is a stationary parametrized 1-cycle, but is not a closed geodesic (and does not correspond to a closed geodesic in any obvious way). This example motivates the following definition: We say that a parametrized 1-cycle $\gamma \in \Gamma_k$ is *strongly stationary* if it is stationary, and for any $i \in \{1, \dots, k\}$ γ_i is a geodesic. If a 1-cycle z is equal to $I(\gamma)$ for some k and some strongly stationary parametrized 1-cycle $\gamma \in \Gamma_k$, then we say that z is a *strongly stationary* non-parametrized 1-cycle. From now on we will be using only the notion of strong stationarity.

It is easy to see that our definition of a non-parametrized strongly stationary 1-cycle is equivalent to the definition of a stationary optimal 1-cycle given in the introduction. Also note that a strongly stationary 1-cycle made of one geodesic segment must be a geodesic loop and therefore must be a closed geodesic. (Two unit vectors tangent to the geodesic loop at its origin must cancel each other.) A strongly stationary 1-cycle made of two geodesic segments either consists of two geodesic segments connecting two different points or consists of two geodesic loops. In the first case it is easy to see that it is a closed geodesic. In the second case we have two subcases. If these geodesic loops are based at different points, then both of them must be closed geodesics. If they are based at the same origin, then we see that the sum of the four unit tangent vectors at the origin of the loops must be equal to zero. If the dimension of the manifold is equal to 2, then this condition implies that this strongly stationary 1-cycle is just a self-intersecting closed geodesic. If the dimension of the manifold is greater than two, then, in principle, these two

geodesic loops need not form a closed geodesic. But it is easy to see that 1) the angles between the tangent vectors to these loops at their common origin p are equal; 2) The bisectors of these two angles in $T_p M^n$ are oppositely directed rays. This condition is, in fact, equivalent to the strong stationarity for a 1-cycle formed by two geodesic loops emanating from the same point.

3. A Morse-theoretic type lemma for Γ_k

The main technical results of this section resemble Theorem 4.3 in [P] (though they do not directly follow from it). On the other hand, they resemble a basic result from the Morse theory asserting that if there are no critical points of a smooth function $F : M \rightarrow R$ on a compact manifold M in the set $F^{-1}([x_1, x_2])$ then the sublevel set $F^{-1}((-\infty, x_1])$ is a deformation retract of the sublevel set $F^{-1}((-\infty, x_2])$. (The deformation retraction can be obtained using the gradient flow of F .) Our goal is to obtain a result of such type for the length functional on Γ_k . The main technical problem is that Γ_k is not an infinite-dimensional manifold, but consists of finitely many intersecting pieces (each of which is an infinite-dimensional manifold).

Lemma 3. *Assume that there are no non-trivial strongly stationary 1-cycles on M^n of length $\leq x$ made of k geodesic segments. Let $f : S^i \rightarrow \Gamma_k^x$ be a continuous map. Then there exists a homotopy $H : S^i \times [0, 1] \rightarrow \Gamma_k^x$, $H(x, 0) = f(x)$, $H(x, 1) \in \Gamma_k^0 \subset \Gamma_k^x$ between f and a map g of S^i into Γ_k such that for any $x \in S^i$ $g(x)$ consists of k constant maps of $[0, 1]$ into M^n .*

Observe that the conclusion of Lemma 3 falls short of the desired contractibility of f . However, if we project everything to the space of non-parametrized 1-cycles, then the contractibility is immediate. Also, we can prove the contractibility if some auxiliary conditions on f are satisfied. More precisely, we have the following proposition:

Proposition 4. *Assume that there are no non-trivial strongly stationary 1-cycles on M^n of length $\leq x$ made of k geodesic segments. Let $f : S^i \rightarrow Z_{(k)}^x$ be a continuous map such that there exists a map homotopic to f that can be lifted to a map $F : S^i \rightarrow \Gamma_k^x$. (That is, f is homotopic to the composition of I and F .) Then f is contractible (in $Z_{(k)}^x$). Further, consider the map $F_j : S^i \rightarrow M^n$ for*

every $j \in \{1, \dots, k\}$ defined by the formula $F_j(p) = (F(p))_j(0.5)$ for any $p \in S^i$. If all these maps F_j are contractible, then F is contractible.

Proof of Proposition 4 assuming Lemma 3: Without any loss of generality we can assume that f is the composition of I and F . Apply Lemma 3 to F . There exists a homotopy H between F and a map of S^i into Γ_k^0 . Observe that I sends all elements of Γ_k^0 to the zero 1-cycle. Therefore the composition of I and H constitutes a homotopy between f and the constant map of S^i into the zero 1-cycle, i.e. a contraction of f .

It remains to verify that if F_j is contractible for each j , then $H(1) : S^i \longrightarrow \Gamma_k^0$ is contractible. Note that $H(1)$ can be regarded as a map of S^i into $(M^n)^k$. Therefore it is sufficient to check only that k maps $H(1)_j$ are contractible. But each of these maps can be connected with F_j by a homotopy Q that can be defined by the formula $Q(t)(p) = (H(t)(p))_j(0.5)$ for any $t \in [0, 1]$, $p \in S^i$. QED.

Proof of Lemma 3:

For the convenience of the reader the proof of Lemms 3 will be split into several steps.

3.1. We are going to demonstrate the existence of a deformation of Γ_k^x into Γ_k^0 , which clearly implies the lemma. Recall that a deformation of a space X into its subset A is, by definition, a continuous map g of X into itself homotopic to the identity map $X \longrightarrow X$ such that $g(X) \subset A$. More generally, if B is a subset of X such that $A \subset B$ we say that a map $g : B \longrightarrow A$ is a deformation in X if there exists a homotopy $G : B \times [0, 1] \longrightarrow X$ such that for any $b \in B$ $G(b, 0) = b$ and $G(b, 1) = g(b)$. But sometimes by a deformation we will mean the whole homotopy G and not just $g(b) = G(b, 1)$.

3.2. Birkhoff curve-shortening process for Γ_k^x . First, we are going to proceed as in the first step of the Birkhoff curve shortening process described in [C] or [ClCo]: Let $\text{inj}(M^n)$ denote the injectivity radius of M^n . Choose $N = \lceil 4x/\text{inj}(M^n) \rceil + 1$. Let γ be an element of Γ_k^x . Divide each of k segments γ_i of γ into N pieces of equal length by points $\gamma_i(t_{ij}), j \in \{0, 1, \dots, N\}, t_{i0} = 0, t_{iN} = 1$. Consider the unique minimizing geodesic segments between $\gamma_i(t_{ij})$ and $\gamma(t_{i,j+1})$ for all j . The length of each of these segments does not exceed $\text{inj}(M^n)/4$. For any $i \leq N$ such geodesic segments form a piecewise geodesic $\bar{\gamma}_i$ connecting $\bar{\gamma}_i(0) = \gamma_i(0)$ with $\bar{\gamma}_i(1) = \gamma_i(1)$. The length of $\bar{\gamma}_i$ does not exceed the length of γ_i . There exists

the following homotopy between γ_i and $\bar{\gamma}_i$: At the moment of time $\tau \in [0, 1]$ we follow each of the N segments of γ_i (between $\gamma_i(t_{ij})$ and $\gamma_i(t_{i,j+1})$) from $\gamma_i(t_{ij})$ to $\gamma_i(\tau t_{ij} + (1-\tau)t_{i,j+1})$ and then make a shortcut from $\gamma_i(\tau t_{ij} + (1-\tau)t_{i,j+1})$ to $\gamma_i(t_{i,j+1})$ along the shortest geodesic. Then we reparametrize the resulting curve proportionally to its arc length. It is easy to see that the length of the curve does not increase during this homotopy. Denote the resulting homotopy by h_i . Combining all of these k homotopies h_i we obtain a homotopy between the parametrized 1-cycle γ and the parametrized 1-cycle $\bar{\gamma} = (\bar{\gamma}_1, \dots, \bar{\gamma}_k)$ made of k piecewise geodesics with N breaks. This homotopy depends on γ in a continuous way. Therefore we obtain a deformation of Γ_k^x into its subset $g_{k,N}^x$ defined as the set of all parametrized 1-cycles $\bar{\gamma} = (\bar{\gamma}_1, \dots, \bar{\gamma}_k)$ from Γ_k^x such that for any i $\bar{\gamma}_i$ is a piecewise geodesic made of N geodesic segments of non-zero length $\leq \text{inj}(M^n)/4$ parametrized proportionally to its arclength. Let us denote this deformation by B_N . Below we will refer to it as the *Birkhoff deformation*. We regard $g_{k,N}^x$ as the subset of a larger set $G_{k,N}^x$ defined as the set of all elements $\gamma = (\gamma_1, \dots, \gamma_k)$ of Γ_k^x such that for any i γ_i is a piecewise geodesic made of at most N geodesic segments of non-zero length $\leq \text{inj}(M^n)/2$. In other words, the only difference between $g_{k,N}^x$ and $G_{k,N}^x$ is that we allow elements of $G_{k,N}^x$ to have somewhat longer geodesic segments.

Now we are going to prove that $g_{k,N}^x$ can be deformed into its subset $g_{k,N}^0 = \Gamma_k^0$ inside Γ_k^x .

3.3 A classification of elements of Γ_k^x and $G_{k,x}^N$. Let us define an equivalence relation on Γ_k^x . For any element $\gamma = (\gamma_1, \dots, \gamma_k)$ of Γ_k^x or $G_{k,N}^x$ consider its $2k$ endpoints $\gamma_i(0), \gamma_i(1)$. We will call these points *multiple points* of γ . The set of these $2k$ points can be partitioned into J non-empty sets A_j , ($J \in \{1, \dots, k\}$), such that 1) Each set A_j contains the equal number of points of the form $\gamma_i(0)$ (for some i) and $\gamma_l(1)$ (for some l); 2) $\gamma_i(t_1) = \gamma_l(t_2)$ for some $i, l \in \{1, \dots, k\}$ and $t_1, t_2 \in \{0, 1\}$ if and only if $\gamma_i(t_1)$ and $\gamma_l(t_2)$ are in the same set A_j for some $j \in \{1, \dots, J\}$. The number J will be called *the number of multiple points* of γ . We will say that two 1-cycles γ and β from Γ_k^x are of the same *type* if these partitions for γ and β coincide, and the set of all i such that γ_i is constant coincides with the set of values of i such that β_i is constant.

For example, let $k = 2$. In these case there will be three types of parametrized

1-cycles from Γ_k^x when neither γ_1 nor γ_2 is constant: (a) $\gamma_1(0) = \gamma_1(1) \neq \gamma_2(0) = \gamma_2(1)$ (1-cycle that consists of 2 closed curves that do not intersect at their endpoints); (b) $\gamma_1(0) = \gamma_2(1) \neq \gamma_2(0) = \gamma_1(1)$ (1-cycles that consist of one closed curve obtained by glueing together γ_1 and γ_2 , where endpoints of γ_1 (and of γ_2) are different; and (c) $\gamma_1(0) = \gamma_1(1) = \gamma_2(0) = \gamma_2(1)$ (γ_1 and γ_2 are loops with the common endpoint. Such 1-cycle can be considered as either made of two closed curves or of one closed curve with the self-intersection.) There will be two types when both γ_1 and γ_2 are constant, namely, $\gamma_1 \neq \gamma_2$ and $\gamma_1 = \gamma_2$. For each $i = 1, 2$ there will also be two types corresponding to the case when γ_i is constant and γ_{3-i} is a non-constant loop: One type corresponds to the case when $\gamma_i(t) = \gamma_{3-i}(0) = \gamma_{3-i}(1)$ and the other type corresponds to the case $\gamma_i(t) \neq \gamma_{3-i}(0) = \gamma_{3-i}(1)$.

However, we will also need a stronger equivalence relation on $G_{k,N}^x \subset \Gamma_k^x$. For each $\gamma \in G_{k,N}^x$ we will consider kN geodesic segments γ_{ij} in k parametrized curves γ_i forming γ . Consider $(N-1)k$ endpoints of these geodesic segments that are not endpoints of k curves γ_i . We will call them *double points* in order to emphasize that there are exactly two geodesic segments meeting at any of these points. (However note, that it is possible that precisely two geodesic segments meet at some of the multiple points as well). We will say that two elements α, β of $G_{k,N}^x$ are of the same *type* as elements of $G_{k,N}^x$ if they are of the same type as elements of Γ_k^x and for each $i = 1, \dots, k$, $j = 1, \dots, N$ the geodesic segment α_{ij} is constant if and only if β_{ij} is constant. Also note that the type of γ as an element of $G_{k,N}^x$ and the (vectors of) positions of J multiple points and $(N-1)k$ double points on the manifold determine an element of $G_{k,x}^N$ uniquely. Therefore for any specific type of γ we can identify an element $\gamma = (\gamma_i)_{i=1}^k$ of $G_{k,N}^x$ with the corresponding element of $(M^n)^{J+(N-1)k}$, where J is the number of multiple points of γ . Note that copies of $(M^n)^{J+(N-1)k}$ corresponding to individual types can be simultaneously embedded into the ambient manifold $(M^n)^{(k+1)N}$ by the corresponding diagonal embeddings that identify endpoints of k intervals in accordance with the type.

3.3.A A partial order on types of elements of $G_{k,N}^x$. We will say that a type A is *higher* than a type B , if: 1) Multiple points of A can be obtained by merging some of the multiple points of B . (In other words the partition considered in 3.3 for A is coarser than the partition for B .); 2) If for some 1-cycle β of type

B , for $i = 1, 2, \dots, k$, and for some $j = 1, \dots, N$ the geodesic segment β_{ij} is constant, then for any element γ of type A γ_{ij} is also constant.

The resulting relation is a partial order on the set of types. For example, the maximal types among types that correspond to cycles of non-zero length are the types with just one multiple point m , $Nk - 2$ constant geodesic segments, and just two non-constant geodesic segments. These two segments correspond to the minimal geodesic connecting m with the only double point, d , and traverse this geodesic in opposite directions.

3.4. Some general remarks about deformations of $g_{k,N}^x$ that will be constructed below. These deformations will consist of finitely many steps. Each of these steps constitute either the Birkhoff deformation described in 3.2, or will be a deformation inside $G_{k,N}^x$. In the last case in order to describe the deformation we need to describe the trajectories of individual multiple and double points. More formally, during a deformation of the last type we will be deforming each individual cycle γ as an element of $(M^n)^{(N+1)k}$. Each multiple or double point will move along trajectories obtained as a projection of trajectories of a vector field defined in an open set in $(M^n)^{(N+1)k}$ that contains the union of all diagonally embedded copies of $(M^n)^{J+(N-1)k}$ corresponding to various types of elements of $G_{k,N}^x$.

We will construct this vector field in such a way that will not allow the type of the elements of $G_{k,N}^x$ to change during these stages of the deformation, but at the very end of the deformation, when the trajectory of the flow hits $G_{k,N}^0$, and the length of the element becomes zero. This is achieved by defining the components of the vector fields that correspond to the individual multiple and double points of γ so that in the situation when the distance between two distinct multiple and/or double points becomes small, they will move along the trajectories of the same smooth vector field on M^n . (Here we are talking only about pairs of points the collision of which can change the type.) Moreover, since the type of any 1-cycle from $g_{k,N}^x$ regarded as an element from Γ_k^x cannot change during the Birkhoff deformation (since all multiple points remain unchanged), Γ_k^x types of 1-cycles remain unchanged through the whole deformation until its very last moment. We will not need this feature of our construction in the proof of Lemma 3, but it will turn out to be convenient for the next section.

However, we can encounter the following problem: We need to ensure that the

distance between any two (double or multiple) points that should be connected by a geodesic segment is less than $\text{inj}(M^n)$. We will resolve this technical complication in the following way. Our choice of N ensures that at the beginning these distances do not exceed $\text{inj}(M^n)/4$ for any $\gamma \in G_{k,N}^x$. Therefore we can deform our parametrized 1-cycles using a flow that will be constructed below for a certain safe amount of time which is sufficiently small to ensure that lengths of segments grow by not more than $\text{inj}(M^n)/4$. Then we stop and apply the Birkhoff deformation with N breaks B_N again. That is, we forget that the k curves forming an element of $G_{k,N}^x$ are already piecewise geodesics with N breaks, divide them into N equal pieces (of length $\leq x/N \leq \text{inj}(M^n)/4$) by $N-1$ points and replace the curve by another piecewise geodesic consisting of N geodesic segments of length $\leq \text{inj}(M^n)/4$ with vertices in these points. (Of course, this replacement is made using the length non-increasing homotopy described in 3.2 that was used in the definition of the Birkhoff deformation.) At this point, in principle, the type of γ as an element of $G_{k,N}^x$ can change because new double points can merge in a different pattern from what was before the Birkhoff stage. (Note however that the Γ_k^x type remain unchanged since B_N does not affect multiple points.) Now we are ready to continue the deformation using the same flow again, etc... The vector fields and the times of these deformations will be chosen so that at each stage of the deformation every element of $G_{k,N}^x$ will become shorter by at least a certain $\delta = \delta(M^n, N, k, x) > 0$. Therefore we will need only a finite number of stages to reach $G_{k,N}^0$.

3.5. The direction of the steepest descent. We will start the construction of the flow from the following observation: Since there is no strongly stationary 1-cycle of positive length $\leq x$, then for any piecewise geodesic 1-cycle γ from $G_{k,N}^x$ there exists a system of vectors at all multiple and double points such that a small deformation of γ (regarded as an element of $(M^n)^{J+(N-1)k}$) in the direction of these vectors leads to an element of $G_{k,N}^x$ of the same type but of a smaller length. These vectors are constructed as follows: Any multiple point corresponds to a set A_j of the partition. The vector at this point is calculated as $\sum_{\gamma_l(t_i) \in A_j; t_i=0 \text{ or } 1} v_l(t_i)$, where $v_l(t_i)$ is the unit vector tangent to γ_l at t_i directed from the multiple point. If we will regard all Nk geodesic segments as curves in M^n , then each double point is adjacent to two geodesic segments of the curve (with the exception

of the case, when this double point is connected by a sequence of geodesic segments of zero length with a multiple point. In this last case the double point coincides with the multiple point, and we *define* the component of v corresponding to this double point to be equal to the already determined component of v corresponding to the multiple point.) The component of v at this double point is calculated as the sum of two unit vectors tangent to two geodesic segments that meet at this point and are directed from it. Our assertion now follows directly from the first variation formula for the length functional. We will call the system of $J + (N - 1)k$ tangent vectors of M^n a *deformation vector* for γ and will denote it by $v(\gamma)$. Note that $v(\gamma)$ is the collection of zero vectors if and only if γ is a strongly stationary 1-cycle.

The first variation of the length of γ in the direction of $v(\gamma)$ is equal to $-\|v(\gamma)\|^2$, where $\|v(\gamma)\|^2$ is calculated as follows: When we deal with J components of v corresponding to multiple points we just sum their squares. We are going to say that a pair of double points *merges* if they are connected by a sequence of *constant* geodesic segments. We define a *cluster* of double points as a maximal set of double points such that each pair of them merges. We say that a cluster is *negligible* if one of double points in the cluster is connected by a sequence of *constant* segments with a multiple point (so geometrically, all double points in this cluster coincide with the multiple point). Our definition of v implies that components of v corresponding to double points in a cluster point are equal. When we calculate $\|v(\gamma)\|^2$ we *by definition* disregard all double points in each negligible cluster and count the squared norm of the component of v corresponding to all double points in a cluster only once for each non-negligible cluster. In other words we define $\|v(\gamma)\|^2$ as $\sum_{m_i; i=1, \dots, J} \|v(\gamma)(m_i)\|^2 + \sum_{\text{Non-negligible clusters of double points}} \|v(\gamma)(d_i)\|^2$, where the first summation is over the set of all multiple points and the second summation is over the set of all non-negligible clusters of double points.

3.6. The deformation vector for γ can be used to decrease the length for all γ_* sufficiently close to γ . Unfortunately, the dependence of $v(\gamma)$ on γ is not continuous. This happens because the type of element of $G_{k,N}^x$ changes in a discontinuous manner. Yet, it is easy to see that for any $\gamma \in G_{k,N}^x$ there exists a sufficiently small open neighborhood U of γ in $G_{k,N}^x$ and a positive μ such that for any $\gamma_* \in U$ a sufficiently small deformation of γ_* in the direction of the deformation vector $T_{\gamma_*}^{\gamma}(v(\gamma))$ defined below decreases the length of γ_* , and the

first variation of the length in the direction of $T_{\gamma}^{\gamma^*}(v(\gamma))$ is less than or equal to $-\|v(\gamma)\|^2/2$. The above deformation vector, $T_{\gamma}^{\gamma^*}(v(\gamma))$ is obtained from $v(\gamma)$ by the parallel transport of all $J + \sum_i (N_i - 1)$ vector components of $v(\gamma)$ along the shortest geodesics connecting the vertices of γ with the corresponding vertices of γ_* .

More precisely, one first chooses U so small that: 1) different multiple or double points of γ cannot merge in U ; 2) For any $\gamma_* \in U$ any of its multiple points has the unique closest multiple point of γ , and any double point of γ_{*i} has the unique closest double or multiple point of γ_i at the distance not exceeding $\text{inj}(M^n)/4$. But note that, in principle, each multiple point of γ can split into two multiple points (that can be connected or not connected by a geodesic segment) or into a pair multiple point - double point for an arbitrarily small U . Also note that if $N_i < N$ then each double point of γ_i can bifurcate into two distinct double points connected by a very short geodesic. If one of k segments γ_i is a constant geodesic loop, then the corresponding multiple point can bifurcate into a pair of points that consists of the multiple point and a double point that is very close to the multiple point. These points are connected by two oppositely oriented copies of the shortest geodesic, together forming a short piece-wise geodesic loop based at the multiple point. Finally note that a finite number of bifurcations of these types can occur simultaneously. So, the dimension of $T_{\gamma}^{\gamma^*}(v(\gamma))$ can be greater than the dimension of $v(\gamma)$. But condition 2) in the definition of U implies that even if a multiple or a double point of γ bifurcates into a finite number of (multiple and/or double) points we know unambiguously how to define the corresponding component of $T_{\gamma}^{\gamma^*}(v(\gamma))$ for each of them: we just perform the parallel transport of the corresponding component of $v(\gamma)$ along the (unique) shortest geodesic.

Now the assertion immediately follows from the continuity on U of the first variation of the length in the direction of the field $X(\gamma_*) = T_{\gamma}^{\gamma^*}(v(\gamma))$. This continuity follows from the first variation formula for the length functional. One just needs to perform easy calculations verifying this continuity for all cases of elementary mergers. That is, it is necessary to consider the particular cases, when $\gamma_i \longrightarrow \gamma$, where all γ_i are of the same type, which can be obtained from (the type of) γ by either 1) a splitting of a multiple point into two multiple points (connected or not connected by a very short geodesic segment converging to the point) or into

the pair “multiple point-double point” (on one of the segments adjacent to the multiple point); or 2) a splitting of a double point into two double points connected by a very short geodesic (converging to the double point). The general case of this formula follows by induction. We will omit the details of this straightforward verification.

3.7. After these preliminaries we are going to prove that:

- A. There exists a sufficiently small positive $\tau_* \leq x$ such that $g_{k,N}^{\tau_*}$ can be deformed to Γ_k^0 ; and
- B. For any positive $\tau \leq x$ there exists a deformation of $g_{k,N}^x$ to $g_{k,N}^\tau$ inside $G_{k,N}^x$.

In fact, as it was already noted, we are going to construct the deformation by first defining a vector flow on an open subset of $(M^n)^{(N+1)k}$, that includes the image of $G_{N,k}^x$ under the embedding discussed above. This flow will be the same for both A and B. So the division of our deformation into these two parts is somewhat artificial. Yet we encounter different difficulties in these two situations: When the length is small, our main problem will be the lack of compactness at zero length of the space $G_{N,k}^x \setminus G_{N,k}^0$, and we need to prove that our flow decreases the length with a speed bounded from zero by a constant. When the length is large, we do not want the distances between double points that must be connected by a geodesic segment, to become large, so from time to time we stop and perform the Birkhoff deformation.

3.8. In order to prove A we would first like to establish a positive lower bound for $\|v(\gamma)\|$ for all $\gamma \in G_{k,N}^{\tau_0} \setminus G_{k,N}^0$ for a sufficiently small positive τ_0 . The key idea is to observe that this statement will be true for the Euclidean space R^n instead of M^n : Assume that there exists a sequence of 1-cycles γ_i in R^n made of at most Nk straight line segments such that $\|v(\gamma_i)\| \rightarrow 0$. Rescale γ_i in R^n so that the maximal length of an edge equals to one. (This does not affect $\|v(\gamma_i)\|$.) Choose a convergent subsequence. Then its limit must be a non-trivial stationary 1-cycle in R^n of length ≥ 1 . (Here we must check what happens with $\|v\|$ when an edge collapses to a point in the limit. It is easy to see that $\|v(\lim_{i \rightarrow \infty} \gamma_i)\| \leq 2 \limsup_{i \rightarrow \infty} \|v(\gamma_i)\|$ in the situation, when we have a sequence of γ_i of the same type, and exactly one segment of γ_i collapses to a point in the limit. The number of such collapses is bounded from above by $Nk - 1$. Therefore the norm of the deformation vector of the limit 1-cycle will be zero.) But

it is very easy to see that there are no stationary 1-cycles in R^n . So, we obtain a contradiction thereby proving the existence of a uniform positive lower bound for $\|v(\gamma)\|$ for all parametrized 1-cycles that consists of at most Nk straight line segments in R^n .

If $\tau_0 = \tau_0(M^n, N, k)$ is sufficiently small, then any parametrized 1-cycle from $G_{k,N}^{\tau_0}$ splits into several connected components contained in very small balls in M^n . Applying the inverse of the exponential map we obtain “almost” a 1-cycle in the tangent space to M^n with “almost” the same angles. Now the existence of a uniform positive lower bound for the norm of the deformation vectors of elements of $G_{k,N}$ for R^n implies the existence of such uniform lower bound for all 1-cycles from $G_{k,N}^{\tau_0}$.

3.9. Construction of a deformation of $g_{k,N}^{\tau_0}$ into Γ_k^0 . Now it is easy to find a countable set $\{\gamma_l\} \subset G_{k,N}^{\tau_0} \setminus \Gamma_k^0$, a locally finite covering of $G_{k,N}^{\tau_0} \setminus \Gamma_k^0$ by open balls U_l centered at γ_l and a subordinate partition of unity that can be used to obtain a continuous function ϕ assigning to every element γ of $G_{k,N}^{\tau_0} \setminus \Gamma_k^0$ a system of tangent vectors to M^n at each of its multiple or double points such that the variation of length of the cycles in direction of $\phi(\gamma)$ is bounded from below by a positive constant δ . (In order to obtain a formal proof of the last assertion we just need to establish the continuity of the first variation of the length. This will be done in 3.10.)

We construct this γ_l and U_l inductively with respect to the partial order on types of elements of $G_{k,N}^x$ introduced in section 3.3.A above. We start from elements of $G_{k,N}^x \setminus G_{k,N}^0$ of the highest possible type that form strata of a high codimension that are closed in $G_{k,N}^x \setminus G_{k,N}^0$. Construct a locally finite covering (in $G_{k,N}^x$) of the union of these strata, so that all centers of open balls forming the covering are on the considered strata. Automatically an open neighborhood of the union of these strata will be covered. Then we proceed to strata corresponding to types of the second highest order. Points of closure of the union of these strata in $G_{k,N}^x \setminus G_{k,N}^0$ that are not in these strata are in strata corresponding to the highest type, and were already covered. Therefore we can choose the covering so that no open ball from this covering intersects an open neighborhood of strata corresponding to the highest types that were previously covered. Also the centers of all balls should be on the strata that are being covered. We continue inductively

in this way until we cover the whole $G_{k,N}^x \setminus G_{k,N}^0$. On each step of the inductive procedure we consider the union strata corresponding to the maximal types that were not yet covered. A neighborhood of the union of all strata corresponding to higher types was already covered on previous steps of the inductive procedure. So we complete the covering of the union of strata that are being considered on the current step by adding open balls centered at the considered strata that have empty intersections with a neighborhood of the union of all already covered strata corresponding to all higher types.

After the covering is completed, and a subordinate partition of unity is chosen, we define $\phi(\gamma)$ as the weighted sum of $T_{\gamma_i}^\gamma(v(\gamma_i))$ over the set of indices i such that $\gamma \in U_i$. (Recall that components of $T_{\gamma_i}^\gamma(v(\gamma_i))$ are obtained by the parallel transport of the corresponding components of $v(\gamma_i)$ along the shortest geodesics between corresponding multiple or double points of γ_i and γ . Of course, U_i should be sufficiently small in order for this definition to be unambiguous, as we explained above.) The weights are equal to the corresponding functions from the partition of unity. This construction provides us with the flow Φ_t that deforms $g_{k,N}^\tau$ to Γ_k^0 in a finite time for each sufficiently small τ (as we will see below).

3.10. The type cannot change during this deformation; the first variation of the length is continuous. Let $p_1, p_2 \in \gamma \in G_{k,N}^x \setminus G_{k,N}^0$ be either two multiple points, or a multiple point and a double point connected by a segment, or two double points connected by a segment. Note that if the distance between p_1, p_2 of γ is very small, then γ is in a small neighborhood of a stratum corresponding to a higher type of cycles, where p_1 and p_2 merge into one point p . If this neighborhood is sufficiently small, then *all* balls of the covering that cover γ are centered at strata corresponding to higher types, where p_1 and p_2 are merged into one point p . But then p_1 and p_2 will be deformed (for some period of time) using the same vector fields on M^n . This will be happening all the time while they will be sufficiently close to each other. But since different integral trajectories of a smooth vector field do not intersect, p_1 and p_2 cannot merge at least until the moment of the deformation, when the total length of the 1-cycle becomes zero.

Also, observe that in the considered situation (when γ is close to a stratum corresponding to a higher type) the first variation of the length in the direction of the vector field that we constructed will be equal to a linear combination of

variations in the direction of vector fields of the form $T_{\gamma_l}^\gamma(v(\gamma_l))$, considered in 3.6. (Here γ_l are located in strata corresponding to higher types, where p_1 and p_2 merge into one point. The coefficients in the linear combination will be the corresponding functions from the partition of unity.) Therefore we can prove the continuity of the first variation of length with respect to γ as in 3.6.

3.11 Note that for any $\gamma \in G_{k,N}^\tau$ $\Phi_t(\gamma)$ is defined only until the moment of time $t(\gamma)$, when the length of $\Phi_t(\gamma)$ will become zero. But one can extend the domain of definition of Φ_t by defining $\Phi_t(\gamma) = \Phi_{t(\gamma)}(\gamma)$ for $t > t(\gamma)$. More precisely, we take $\tau_* = \min\{x, \tau_0, \delta, \text{inj}(M^n)/4\}$, where τ_0 is as in 3.8, and δ is the lower bound of the speed of decrease of the length introduced at the beginning of 3.9, and just follow the flow until we hit Γ_k^0 . It is clear that: 1) For any element $\gamma \in g_{k,N}^\tau$ we will reach Γ_k^0 in time $t(\gamma) \leq 1$; 2) Since the total length of γ decreases, the distance between any two points on M^n that should be connected by the shortest geodesic in order to obtain $\Phi_t(\gamma)$ does not exceed $\text{inj}(M^n)/4$. Therefore $\Phi_t(\gamma)$ is unambiguously defined. (Recall that we move multiple and double points of γ along trajectories of vector fields determined by the corresponding components of $\phi(\gamma)$. In order to obtain $\Phi_t(\gamma)$ we connect these points by the shortest geodesics.) 3) $t(\gamma)$ depends on γ continuously (by virtue of the implicit function theorem. In order to apply the implicit function theorem we need to establish the continuous differentiability of the length as a function of γ , but it is equivalent to the continuity of the first variation of the length of γ in the direction of the vector field that was established in the previous subsection.)

Therefore the map assigning to γ the point $\Phi_{t(\gamma)}(\gamma) \in \Gamma_k^0$, where the trajectory of the flow reaches Γ_k^0 is continuous, and is the deformation of $G_{k,N}^\tau$ to Γ_k^0 .

3.12. It remains to prove the existence of a deformation of $g_{k,N}^x$ to $g_{k,N}^{\tau_*}$ inside Γ_k^x . Use the compactness of the closure S of $G_{k,N}^x \setminus G_{k,N}^{\tau_*/2}$ to find a finite open covering of S by open neighborhoods U_l of $\gamma_l \in S$ such that for any $\gamma_* \in U_l$ the first variation of the length of γ in the direction of $T_{\gamma_l}^{\gamma_*}(v(\gamma_l))$ does not exceed $-\|v(\gamma_l)\|^2/2$. (Recall that we have proved the existence of an open neighborhood U with this property for any $\gamma \in G_{k,N}^x \setminus \Gamma_k^0$.) Using a subordinate partition of unity $\alpha_l(\gamma)$ define a vector field $\phi(\gamma)$ by the formula $\phi(\gamma) = \sum_l \alpha_l(\gamma) T_{\gamma_l}^\gamma(v(\gamma_l))$, where we perform the summation only over indices l such that $\gamma \in U_l$.

Here the construction of this open covering is similar to that in section 3.9: We

construct it inductively starting from the strata that correspond to the highest type and then proceed to cover strata corresponding to lower types. On each step we add only open balls centered at points on the considered strata that do *not* intersect with an open neighborhood of the already covered strata corresponding to higher types. As the result the $G_{k,N}^x$ type will not be changing during the considered stage of the deformation, and the proof of this fact coincides with the proof in section 3.10 above almost verbatim.

Rescale $\phi(\gamma)$ by a continuous function equal to zero on $G_{k,N}^{\tau_*/2}$ and to one on $G_{k,N}^{\tau_*}$. Denote the resulting vector field by $\psi(\gamma)$. Let $t_* = \text{inj}(M^n)/(16k)$. Consider the flow $\Psi_t(\gamma)$ defined for all $\gamma \in g_{k,N}^x$ at $t \in [0, t_*]$ and determined by the vector field $\psi(\gamma)$. Our choice of t_* guarantees that $\Psi_t(\gamma)$ will be in $G_{k,N}^x$. (In other words, the distance between any pair of points that need to be connected by a geodesic segment will not exceed $\text{inj}(M^n)/2$.) Observe that for any $\gamma \in g_{k,N}^x \setminus G_{k,N}^{\tau_*}$ the difference between the length of γ and the length of $\Psi_{t_*}(\gamma)$ will be at least δt_* , where $\delta = \frac{1}{2} \min_{l=1}^l \|v(\gamma_l)\|^2$.

Now recall that the Birkhoff curve shortening process provides us with the deformation B_N of Γ_k^x into $g_{k,N}^x$. The restriction of B_N to $G_{k,N}^x \subset \Gamma_k^x$ is a deformation of $G_{k,N}^x$ into $g_{k,N}^x$ inside Γ_k^x . Apply B_N . The composition of B_N and Ψ_{t_*} is a curve-shortening deformation of $g_{k,N}^x$ into $g_{k,N}^{\max\{x-\delta t_*, \tau_*\}}$ inside Γ_k^x .

Now we can apply Ψ_{t_*} and then B_N again and again, etc. Let $K = \lceil \frac{1}{\delta t_*} \rceil + 1$. It is easy to see that $(B_N \Psi_{t_*})^K$ is the required deformation of $g_{k,N}^x$ to $g_{k,N}^{\tau_*}$ inside Γ_k^x . QED.

Observation. We would like to mention again (for a further use) that the defined in 3.3 type of elements of Γ_k^x does not change during the considered deformation until possibly the very last moment, when the length becomes zero.

Remark. It seems that one can find a more straightforward proof of a modification of Proposition 4 suitable for our purposes using the same approach as in the proof of Theorem 4.3 in [P] (but we had not checked the details). The idea is to use the space of varifolds, the compactness of closed balls in the space of varifolds (that follows from the Banach-Alaoglu theorem), and the continuity of the first variation of the mass of a varifold in the direction of a fixed vectors field (with respect to the variable varifold). Then, as in the proof of Lemma 3, one constructs a map from the space of varifolds of the bounded mass to the space of C^1 -smooth vector fields

such that the mass decreases fast, when we apply the corresponding flow. One can use an appropriate locally finite covering of the space of varifolds of bounded non-zero mass and a subordinate partition of unity in order to construct this map to the space of C^1 -smooth vector fields. It seems that in this way one can avoid minor combinatorial complications in our proof related to the combinatorics of the space of parametrized 1-cycles made of piecewise geodesics.

Finally observe that if M^n is diffeomorphic to S^2 we can combine Lemma 3 in the case of $k = 2$ with the observation made in the last paragraph of section 2 to obtain an elementary proof of the following assertion used in our paper [NR]. (This assertion first appeared in [ClCo].)

Proposition 5. *Let M be a Riemannian manifold diffeomorphic to S^2 . Assume that for some x there exists a non-contractible map $f : S^1 \longrightarrow \Gamma_2^x$. Then there exists a non-trivial closed geodesic of length $\leq x$ on M .*

Proof. Lemma 3 implies the existence of a non-trivial strongly stationary 1-cycle in Γ_2^x . But, as we noted at the end of section 2, each strongly stationary 1-cycle on a two-dimensional manifold made of two segments is either a closed geodesic or a union of two closed geodesics. But our strongly stationary 1-cycle is non-trivial. Therefore either it is a non-trivial closed geodesic or it contains a non-trivial closed geodesic as a subset. QED.

4. Almgren correspondence

Now we are going to explain the Almgren correspondence in the 1-dimensional case for 1-cycles made of finitely many closed curves. (This simplified version is all that we will need for our purposes - see [A] for the full story). Assume that we are given a continuous map A of a disk D^m or a sphere S^{m-1} , or more generally, a compact polyhedron $|K|$ into Γ_k . The Almgren correspondence assigns to A a $(\dim|K| + 1)$ -dimensional singular chain on M^n as follows: We can regard Γ_k as a subset of the topological space of all maps of the disjoint union of k copies of $[0, 1]$ into M^n . To any of these maps we can assign a continuous map of $X = D^m \times \bigcup_{i=1}^k [0, 1]_i$ (or $X = S^{m-1} \times \bigcup_{i=1}^k [0, 1]_i$, or $X = |K| \times \bigcup_{i=1}^k [0, 1]_i$) into M^n in the standard way. Further, since elements of Γ_k are parametrized cycles we can identify points of $2k$ sets $D^m \times \{0\}$, $D^m \times \{1\}$ (or, correspondingly $2k$ sets $S^{m-1} \times \{0\}$, $S^{m-1} \times \{1\}$, or $|K| \times \{0\}$, $|K| \times \{1\}$) that are mapped into the

same points of M^n . The resulting quotient X_A of X can be quite complicated. For our purposes we will only need the situation when X_A can be triangulated with a finite number of simplices. Moreover, we are going to make an even stronger assumption that will always hold when we apply the Almgren correspondence: We assume that the compact polyhedron K has a simplicial subdivision such that for any open simplex σ of any dimension of this subdivision all parametrized 1-cycles $A(t)$, $t \in \sigma$ have the same type. We will call this assumption a *local triviality* of A . Now we can triangulate X_A so that the obvious projection $X_A \rightarrow K$ becomes a simplicial map and consider the corresponding singular chain in M^n . The local triviality of A makes the following assertions evident: If $K = S^{m-1}$ then that the resulting singular m -chain will be a singular cycle. Its homology class does not depend on the chosen triangulation of X_A . If A is a map of S^{m-1} to Γ_k obtained as the restriction of a map B of D^m to Γ_k then for any triangulation of X_B the boundary of the corresponding singular $(m+1)$ -chain in M^n will be the singular chain obtained from $X_A \subset X_B$ with the induced triangulation. Therefore the singular m -cycle in M^n assigned to $A : S^{m-1} \rightarrow M^n$ will represent 0 in $H_m(M^n)$ if A is contractible.

Note that if f is a map of S^{i-1} into Γ_k^x satisfying the local triviality assumption, and $H : S^{i-1} \times [0, 1] \rightarrow \Gamma_k^x$ is the homotopy between f and a map g of S^{i-1} into Γ_k^0 constructed as in the proof of Lemma 3 (in the situation when there are no non-trivial strongly stationary parametrized 1-cycles in Γ_k^x), then H also satisfies this assumption. This assertion immediately follows from the observation made right after the proof of Lemma 3. This observation will easily imply the local triviality of A in situations when we will need to use the Almgren correspondence in the course of proving Theorems 1 and 2 in the next section.

Yet for the sake of completeness we are also going to sketch how to modify the above version of the Almgren correspondence in order to make it work in the general case, when we do not even have the triangulability of X_A . (This construction will not be used in the present paper.) Recall that at the beginning of the proof of Lemma 3 in the previous section we introduced $N = N(M^n, x) = [4x/\text{inj}(M^n)] + 1$, the spaces $g_{k,N}^x$ and $G_{k,N}^x$ made of parametrized 1-cycles of length $\leq x$ formed by k piecewise geodesics made of at most N geodesic segments of length $\leq \text{inj}(M^n)/4$ and $\leq \text{inj}(M^n)/2$, correspondingly, parametrized

proportionally to the arclength. We also defined the curve-shortening Birkhoff deformation of Γ_k^x into $g_{k,N}^x$. Further recall that $g_{k,N}^x$ and $G_{k,N}^x$ can be regarded as subsets of M^{kN} . It is easy to prove that there exists a subset $\bar{g}_{k,N}^x$ of M^{kN} containing $g_{k,N}^x$ and contained in $G_{k,N}^x$ that can be triangulated. (The shortest formal way to prove the last assertion is the following: Approximate the Riemannian metric on M^n in C^3 -topology by an analytic Riemannian metric so that the distances on the resulting Riemannian manifold $\bar{M}^n$ do not exceed corresponding distances on M^n . Now observe that metric balls of radius $\leq \text{inj}(M^n)/2$ on $\bar{M}^n$ are subanalytic sets, the restriction of the distance function to such metric balls is a subanalytic function, and that according to a well-known theorem of H. Hironaka subanalytic sets are triangulable (cf. [B] for more details. See also [BM] for the definition and basic properties of subanalytic sets and function, including the proof of the mentioned theorem of H. Hironaka.) Therefore we can define $\bar{g}_{k,N}^x$ as $g_{k,N}^x$ but using the distance function on $\bar{M}^n$ instead of the distance function on M^n .) It is easy to see that one can triangulate $\bar{g}_{k,N}^x$ so that the type of parametrized 1-cycles is constant on every simplex of the triangulation. Therefore we can take $x = \max_{y \in K} l(A(y))$, compose B_N with A , and take a simplicial approximation $\bar{A}$ of the resulting composition $B_N A : K \longrightarrow \bar{g}_{k,N}^x$. Now we can consider the quotient $X_{\bar{A}}$ defined as above. It is easy to see how to triangulate $X_{\bar{A}}$. Therefore we can proceed as above, and consider the corresponding singular chain in M^n . It is clear that if $K = S^{m-1}$, then this chain is a cycle, and this cycle represents $0 \in H_{m-1}(M^n)$ if and only if A is contractible. A small technical complication that arises here is the following: Assume that we apply the Almgren correspondence to a map $B : D^m \longrightarrow \Gamma_k^x$ and to the restriction A of B to $S^{m-1} = \partial D^m$. Then the values of x defined for these two mappings will, in general, be different. Therefore the cycle in M^n corresponding to A will not, in general, coincide with the boundary of the chain corresponding to B . Yet, it is easy to construct a homology between these two cycles.

In the case of a map of a polyhedron into $Z_{(k)}$ one can consider a sufficiently fine subdivision of the polyhedron K . For any simplex of this subdivision the restriction of our map onto this simplex lifts to Γ_k . Then we can proceed as in the parametrized case (for this simplex). Finally sum the resulting chains over all simplices of the triangulation.

As it was noted, F. Almgren described in [A] a similar but somewhat more technically complicated construction applicable to maps of a polyhedron into $\bar{Z}_1(M^n, Z)$ or even $\bar{Z}_k(M^n, Z)$.

5. Proofs of Theorems 1 and 2.

Proof of Theorem 1. Assume that $\alpha(M^n)$ and, in particular, $l(M^n)$ is greater than $\frac{(n+2)!d}{3}$. Consider a map $f : S^q \rightarrow M^n$ representing a non-zero element of $\pi_q(M^n)$, where S^q is the standard sphere with a fine triangulation. Let $[S^q]$ be the fundamental class of S^q . Since the map is non-contractible, $f_*([S^q]) \neq 0 \in H_q(M^n)$. Let D^{q+1} be a disc that has S^q as its boundary. Triangulate D^{q+1} as the cone over the triangulation of S^q (introducing one new 0-dimensional simplex at the centre of D^{q+1}). We will try to construct a singular $(q+1)$ -chain in M^n , such that $f_*[S^q]$ will be its boundary, which is clearly impossible and will result in a desired contradiction.

We are going to proceed inductively assigning an i -dimensional singular chain in M^n to each i -dimensional simplex of D^{q+1} on the i -th step. This assignment will be denoted by F . The boundary of the singular chain that corresponds to an arbitrary simplex σ_i will be equal to the signed sum of chains assigned to simplices of the boundary of σ_i . These signs will be the same as the signs with which the corresponding simplices enter $\partial\sigma_i$. These singular i -chains will be obtained from $(i-1)$ -dimensional discs in the space of 1-cycles, particularly in $Z_{(k(i-1))}$ for some function $k(i)$. One will use the Almgren correspondence between discs and chains explained in Section 4. In turn, these $(i-1)$ -dimensional discs are obtained by contracting $(i-2)$ -dimensional spheres in $Z_{(k(i-2))}$ that are constructed from $(i-2)$ -dimensional discs in $Z_{(k(i-2))}$ corresponding to simplices of $\partial\sigma_i$ that were constructed on the previous stage of induction.

Alternatively, we can describe this procedure in the following equivalent way (with somewhat more details): We start from a collection of maps of $D^1 \rightarrow Z_{(k(1))}$, where $k(1) = 3$, described below. Then, inductively, for each simplex σ^i in the considered triangulation of D^{q+1} , ($i = 2, 3, \dots, q+1$), we do the following: 1) Construct a map of S^{i-2} into $Z_{(k(i-1))}$ using $(i+1)$ maps of D^{i-2} into $Z_{(k(i-2))}$ corresponding to $i+1$ $(i-1)$ -dimensional simplices in the boundary of σ^i and obtained on the previous step of induction;

2) We use our assumption about non-existence of sufficiently short non-trivial strongly stationary 1-cycles and Proposition 4 to obtain a map of D^{i-1} into $Z_{(k(i-1))}$ contracting this map of S^{i-2} . This map of D^{i-1} will correspond to σ^i .

All these maps from discs and spheres into $Z_{(k(i))}$, $i = 1, 2, \dots, q$, can be lifted to $\Gamma_{k(i)}$. (In fact, these maps will be obtained as compositions of maps into $\Gamma_{k(i)}$ and projections $\Gamma_{k(i)} \longrightarrow Z_{(k(i))}$). So, after the completion of this induction process for every $(q+1)$ -dimensional simplex of the considered triangulation of D^{q+1} we obtain a map from D^q to $Z_{(k(q))}$, that can be lifted to $\Gamma_{k(q)}$. Then we will apply the Almgren correspondence to each of the resulting maps of D^q into $\Gamma_{k(q)}$ and sum the resulting singular $(q+1)$ -chains in M^n . The result will be the required $(q+1)$ -chain.

We will begin with the **0-skeleton** of $D^{q+1} \setminus S^q$ that consists of the point p , the center of the disc. We will assign to p a singular 0-chain that corresponds to an arbitrary point $\tilde{p} \in M^n$. Now we will proceed to the **1-skeleton**: we will assign to the 1-simplices of the form $[v_i, p]$ the singular 1-chains that correspond to minimal geodesics in M^n that connect $\tilde{p}$ and $\tilde{v}_i = f(v_i)$. Next, we consider the **2-skeleton**: Let $\sigma^2 = [v_i, v_j, p]$ be a 2-simplex of $D^{q+1} \setminus S^q$. Consider its boundary $\partial\sigma^2$ and the corresponding singular 1-chain on M^n , which equals to $[\tilde{v}_j, \tilde{p}] - [\tilde{v}_i, \tilde{p}] + [\tilde{v}_i, \tilde{v}_j]$. This can be viewed as a curve of length $\leq 2d + \epsilon$. By our assumption, there is no closed geodesics of length smaller than or equal to $2d + \epsilon$, so there is a curve shortening homotopy that connects this curve with a point. Therefore, we assign to this 2-simplex a singular 2-chain consisting of one singular 2-simplex that corresponds the surface generated by this homotopy. The “extension” to the **3-skeleton** will be somewhat different. Let $\sigma^3 = [v_{i_0}, v_{i_1}, v_{i_2}, v_{i_3}]$ be a 3-simplex of $D^{q+1} \setminus S^q$. We want to find a singular 3-chain to assign to this simplex. Consider $\partial\sigma^3$. There is a singular 2-chain assigned to the boundary of this simplex, which can also be viewed as a 2-sphere in M^n of a particular shape. Namely, to each of the faces of the boundary not in S^q there was assigned a surface generated by a curve shortening homotopy. Without any loss of generality we can assume that the chosen fine triangulation of S^q and the map of S^q into M^n were chosen so that any two-dimensional simplex of the triangulation S^q is also mapped into the surface obtained by contracting its boundary in M^n by a

homotopy that does not increase the length. As we will see, this 2-sphere corresponds to a 1-sphere in the space $Z_1(M^n, Z)$ that passes through the subset that consists of cycles that are composed of no more than 12 curves, and the mass of each such cycle is bounded from above by $8d + 4\epsilon$. (See figure 1 to understand how this 1-sphere is constructed.) In order to describe this correspondence let $e_1 = [\tilde{v}_{i_0}, \tilde{v}_{i_1}]$, $e_2 = [\tilde{v}_{i_0}, \tilde{v}_{i_2}]$, $e_3 = [\tilde{v}_{i_0}, \tilde{v}_{i_3}]$, $e_4 = [\tilde{v}_{i_1}, \tilde{v}_{i_2}]$, $e_5 = [\tilde{v}_{i_1}, \tilde{v}_{i_3}]$, $e_6 = [\tilde{v}_{i_2}, \tilde{v}_{i_3}]$, where each $[\tilde{v}_{i_s}, \tilde{v}_{i_t}]$ is a minimal geodesic segment on the manifold. Then we will let $\gamma_1 = e_1 + e_5 - e_3$, $\gamma_2 = -e_1 + e_2 - e_4$, $\gamma_3 = -e_2 + e_3 - e_6$, $\gamma_4 = e_6 - e_5 + e_4$. Let x_i be a point to which γ_i contracts for $i = 1, \dots, 4$. Then the 1-sphere in the space of 1-cycles will be constructed as follows: let $\tilde{f}_i : D^2 \rightarrow M^n$, $i = 1, \dots, 4$ be each of the four discs that make the 2-sphere in M^n . Those discs correspond to four maps $f_i : [0, 1] \rightarrow Z_1(M^n, Z)$, such that $f_i(0) = T_{\{x_i\}} = 0$, $f_i(1) = T_{\gamma_i}$. These maps are precisely curve-shortening homotopies used to obtain $\tilde{f}_i$; for any $t \in [0, 1]$ $f_i(t)$ is a 1-cycle that consists of one closed curve. It can be regarded as an element of $Z_{(3)}$ if we represent γ_i as the collection of three curves (=three sides of the triangle) glued at their endpoints, and will keep track of these three curves during homotopies contracting γ_i . Now we will let $G_1 : [0, 1] \rightarrow Z_{(12)}$ be the map that for each $q \in [0, 1]$ assigns $\sum_{i=1}^4 T_{f_i(q)}$, (see figure 1(b)). Note that $G_1(0) = \sum_{i=0}^4 T_{\{x_i\}}$, which is the zero cycle, (see figure 1(a)) and that $G_1(1) = \sum_{i=0}^4 T_{\gamma_i}$, which is also the zero cycle, (see figure 1(c)). Thus, we obtain a map from S^1 to $Z_1(M^n, Z)$.

Proposition 4 implies that one of the following is true about this 1-sphere: either it can be contracted to a point without the mass increase, or there exists a stationary 1-cycle of order 12 of mass bounded by $8d + 4\epsilon$. The existence of such a cycle for all sufficiently small ϵ is precluded by our assumption. So, there a disc that passes through 1-cycles of order 12 of mass bounded by $8d + 4\epsilon$. In order to apply Proposition 4 here we first must check that our map of S^1 into $Z_{(12)}^{8d+4\epsilon}$ is homotopic to a map that lifts to Γ_{12} . The lifting of the map of $[0, 1] \in S^1 = [0, 1]/0, 1$ to $Z_{(12)}$ is obvious, but we also need to find a homotopy of $\bigcup_{i=1}^4 \gamma_i$ to $\bigcup_{i=1}^4 \{x_i\}$, where each $\{x_i\}$ is counted three times and is regarded as a constant segment, in Γ_{12} . This can be achieved by first cancelling in a continuous way six pairs of edges e_i with the opposite orientations to a point, which is obviously possible (each pair is connected over itself to the point corresponding to $t = \frac{1}{2}$ counted twice), and then connecting 12-tuples of these points regarded as an element of

Γ_{12} with the constant 1-cycle $\{x_1, x_2, x_3, x_4\}$ regarded as the cycle from Γ_{12} (each point is counted three times) using twelve continuous paths. These paths follow our homotopies restricted to the points of edges of γ_i corresponding to $t = 0.5$ in the chosen parametrization of these edges. As the result we obtain a lifting to Γ_{12} of a map that differs from f only by a reparametrization.

Further, the last assertion of Proposition 4 implies that the lifting of our map $S^1 \rightarrow Z_{(12)}^{8d+4\epsilon}$ to Γ_{12} is also contractible. Indeed, we just need to verify the contractibility of 12 maps of $S^1 \rightarrow M^n$. Of course, this fact follows from the simply-connectedness on M^n . However, there is even a more straightforward geometric reason for contractibility of these 12 circles in M^n : each of them is formed by the trajectory of a homotopy $\tilde{f}_i$ from x_i to a point in the middle of a geodesic segment e_j traversed two times in the opposite directions.

Using the Almgren correspondence we see that this disc corresponds to a 3-chain that we will denote $\tilde{C}_{v_{i_0}, \dots, v_{i_3}}$ in M^n that has $F(\partial\sigma^3)$ as its boundary. So, we will assign $\tilde{C}_{v_{i_0}, \dots, v_{i_3}}$ to the simplex σ^3 . Now suppose we want to extend to **4-skeleton**. Consider any 4-simplex of $D^{q+1} \setminus \partial D^{q+1}$ $\sigma^4 = [v_{i_0}, \dots, v_{i_4}]$. The following 3-dimensional cycle in M^n : $C_{v_{i_0}, \dots, v_{i_4}} = \sum_{j=0}^4 (-1)^j \tilde{C}_{v_{i_0}, \dots, \hat{v}_{i_j}, \dots, v_{i_4}}$ corresponds to the boundary $\partial\sigma^4$ of this simplex. We claim that there exists a corresponding map of the 2-disc to $Z_1(M^n, Z)$, that takes the boundary of this disc to the zero cycle, and such that the resulting sphere passes through the subset that consists of cycles of order 60, i.e. cycles that are composed of no more than twenty closed curves (each closed curve being composed of three segments), of total mass no more than $20(2d+\epsilon)$. So, this map represents an element of $\pi_2(Z_1(M^n))$, and this element will be precisely the element corresponding to the homology class of the considered 3-dimensional cycle in M^n under the Almgren isomorphism. The above map, denoted $G_2 : \bar{D}^2 \rightarrow Z_1(M^n, Z)$ will be constructed as follows: let $f_j : \bar{D}^2 \rightarrow Z_1(M^n, Z)$ be a map corresponding to $(-1)^j \tilde{C}_{v_{i_0}, \dots, \hat{v}_{i_j}, \dots, v_{i_4}}$. Then let $G_2(q) = \sum_{j=0}^4 T_{f_j(q)}$ for any $q \in \bar{D}^2$.

Let us now examine $G_2(\partial\bar{D}^2)$. We will see that for any $q \in \partial\bar{D}^2$, $G_2(q)$ will correspond to the union of 10 pairs of closed curves, where each pair will contain the same curve with two different orientations. In other words, the corresponding 1-cycles will have opposite signs, and will cancel. As the result we obtain a zero cycle.

Thus, we obtained a 2-sphere in the space of 1-cycles. We would like to apply Proposition 4. In order to do that we need to lift that map G_2 to the map $\tilde{G}_2 : \bar{D}^2 \rightarrow \Gamma_{60}$ and examine what happens to the boundary of the disc under this map. Each point on the boundary is mapped to the 30 pairs of segments in M^n . Each pair consists of the same segment with opposite orientations. We want to construct a homotopy between $\tilde{G}_2 : \partial\bar{D}^2 \rightarrow \Gamma_{60}$ and a constant map, i.e. a map that will take a circle to 60 point curves. To construct this homotopy we cancel pairs of parametrized 1-cycles corresponding to the 1-cycles with opposite orientations mentioned in the previous paragraph in a continuous way. We contract each pair $\gamma \cup -\gamma$ to $\gamma(0.5)$ over γ . Thus, we obtain a circle in the space Γ_{60} , where each point $p \in S^1$ corresponds to 60 constant paths (those paths are different for $p \neq p'$). This circle can be interpreted as 60 circles on M^n . Since M^n is simply connected, these circles can be contracted to an arbitrary point in M^n . After contracting them we can obtain a point in the space Γ_{60} made of 60 constant segments. So, combining $\tilde{G}_2$ with these two homotopies, we obtain a map of the 2-disc into Γ_{60} such that its boundary is mapped into a point composed of 60 constant segments. We can factor this map through the sphere S^2 obtained from the disc by identifying its boundary to a point (say, the north pole of the sphere. In this case the southern hemisphere is mapped by $\tilde{G}_2$, and the northern hemisphere is mapped into the subset of Γ_{60} that corresponds to the zero cycle (i.e. in $I^{-1}(0)$.) So Proposition 4 applies: We have constructed a 2-dimensional sphere in the space Γ_{60} , and can conclude that either this sphere can be contracted along the cycles of mass $\leq 20(2d + \epsilon)$, or we have a stationary 1-cycle of mass controlled from above by this bound. (Here the verification of the contractibility of maps F_j defined in the text of Proposition 4 is equivalent to the contractibility of certain 60 2-spheres in M^n . But now we are discussing the case of $q \geq 3$, so M^n is 2-connected.) If ϵ is sufficiently small, then the second case is impossible. In the former case, we obtain a 3-disc in the space of 1-cycles, that corresponds to a 4-chain that we will denote $\tilde{C}_{v_0, \dots, v_4}$. We will then assign this chain to σ^4 .

Now we can continue in the above manner until we fill the original q -dimensional chain $f_*([S^q])$ by a $(q + 1)$ -dimensional chain in M^n . As a corollary of our assumption nothing will stop us until we construct the desired filling. But as it was said before, this is impossible, and we obtain a contradiction refuting our

assumption. The constants $(q+2)!/3$ and $(q+2)!/2$ in the text of Theorem 1 can be explained by the fact that all our 1-cycles consist of at most $4 \times 5 \times 6 \times \dots \times (q+2)$ closed curves of length not exceeding $2d+\epsilon$, and each of these closed curves consists of three segments.

Note that we can get a better estimate when $q = 2$. In this case we need to perform the extension process only till the dimension $q + 1 = 3$. We will need “to represent” the union of four maps of D^2 to M^n corresponding to four faces of a 3-dimensional simplex as a map of a circle to $Z_1(M^n, Z)$. (Recall that these four maps were obtained by contracting the maps of boundaries of these discs to a point without increase of their lengths, see Fig. 1). In the body of the proof we mapped a generic point $t \in [0, 1]$ into the 1-cycle that corresponds to the union of four curves obtained from homotopies contracting ∂D_i^2 at the moment t (see Fig. 1(b)). In the particular case $q = 2$ we can proceed in a slightly different way. We can start from two points obtained as the result of contraction of the maps of boundaries of D_1^2 and D_2^2 and to pass via cycles made of two closed curves (obtained during the curve-shortening homotopies contracting the maps of ∂D_1^2 and ∂D_2^2) to the cycle made of the images of these two boundaries (see Fig.1). The edge $[v_0, v_2]$ will be passed twice with opposite orientation. Continue the homotopy by cancelling this edge. At the end of this homotopy we obtain the map of the boundary of $D_1^2 \cup D_2^2$. Now note that $\partial(D_1^2 \cup D_2^2) = \partial(D_3^2 \cup D_4^2)$. But we can similarly construct a homotopy between $\partial(D_3^2 \cup D_4^2)$ and the zero cycle that uses 1-cycles made of two curves obtained from the curve-shortening homotopies contracting the maps of the boundaries of D_3^2 and D_4^2 . Joining these two homotopies we obtain the desired homotopy between the zero 1-cycle and the zero 1-cycle, i.e. the desired circle in the space of 1-cycles that passes through 1-cycles made of not more than two closed curves of length not exceeding $2d + \epsilon$ (each). See the proof of Theorem 1 in [NR] for more details (in the situation when M^n is diffeomorphic to S^2 . But this part of the proof is the same there as in the more general situation.) QED.

Proof of Theorem 2. Assume $\alpha(M^n)$ and, in particular, $l(M^n)$ is greater than $2(n+1)n^n n!^{\frac{1}{2}}(n+2)!(\text{vol} M^n)^{\frac{1}{n}}$. Then $\alpha(M^n)$ (and $l(M^n)$) are greater than $(n+2)! \text{FillRad}(M^n)$. The definition of the filling radius implies that M^n bounds in the $(\text{FillRad}(M^n) + \delta)$ -neighborhood of M^n in $L^\infty(M^n)$. Let W “fill” M^n

in the $(FillRad(M^n) + \delta)$ -neighborhood of M^n (that is $M^n = \partial W \pmod{2}$): Since we do not assume that M^n is orientable, its fundamental homology class $[M^n]$ is defined only for Z_2 coefficients.) Without any loss of generality we can assume that W is a polyhedron.

Suppose W together with M^n is endowed with a very fine triangulation. We are going to try to construct a singular $(n+1)$ -chain on M^n such that the boundary of that chain is homologous to the boundary of W (regarded as a chain). That is clearly impossible, so we will obtain a contradiction. We will construct this chain by induction with respect to the dimension of skeleta of W . That is to each i -simplex of W we will assign a singular i -chain on M^n . We will begin with the **0-skeleton** of W . Let v_i be a vertex of W . Then $F(v_i) = \tilde{v}_i \in M^n = \partial W$, such that $d(v_i, \tilde{v}_i) = d(v_i, M^n) \leq FillRad M^n + \delta$. Suppose $\tilde{v}_i, \tilde{v}_j$ come from the vertices v_i, v_j of some simplex in W . Then $d(\tilde{v}_i, \tilde{v}_j) \leq 2FillRad M^n + 3\delta$. (We assume here that the triangulation of W is fine so that the lengths of 1-simplices of the triangulation are at most δ .) Next, we are going to extend F to the **1-skeleton**. We will assign to any 1-simplex $[v_i, v_j] \subset W \setminus M^n$ a singular 1-chain that corresponds to a minimal geodesic that connects $\tilde{v}_i$ and $\tilde{v}_j$ of length $\leq 2FillRad M^n + 3\delta$. Now we can see that the boundary of each 2-simplex in W is sent to a singular chain that corresponds to a curve of length $\leq 6FillRad M^n + 9\delta$, (we will assume that all simplices in M^n are already short).

Next we are going to extend to the **2-skeleton**. Let σ^2 be a 2-simplex of W . Consider its boundary $\partial\sigma^2$ and its corresponding singular 1-chain. There is a curve shortening homotopy that connects the curve corresponding to that chain to a point. So we will map σ^2 to the chain that corresponds to the surface determined by this homotopy. To “extend” F to the **3-skeleton** of W consider an arbitrary 3-simplex σ^3 . Consider its boundary $\partial\sigma^3$ and the corresponding singular 2-chain, which can be viewed as 1-sphere in the space $Z_1(M^n, Z)$ as in the proof of the Theorem 1. This sphere passes through 1-cycles of length $\leq 4(6FillRad M^n + 9\delta)$. Suppose this sphere cannot be contracted via the 1-cycles of smaller mass. Then there exists minimal 1-cycle of length $\leq 4(6FillRad M^n + 9\delta)$ contradicting our assumption. (Here we use Proposition 4 from the previous section. One can check that our spheres in the space of non-parametrized 1-cycles can be lifted to spaces of parametrized 1-cycles exactly as this was done in the proof of Theorem 1 above.)


So the above 1-sphere can be “filled” by a disc that passes through 1-cycles of mass not exceeding the above bound. This disc corresponds to a singular 3-chain that has $F(\partial\sigma^3)$ as its boundary. So we will assign this chain to σ^3 . The procedure of “extending” to 4-skeleton is similar to the one in the proof of the Theorem 1: At this point “the image” of $\partial\sigma^4$ has been determined and it equals to $C_{v_{i_0}, \dots, v_{i_4}} = (-1)^j \sum_{j=0}^4 \tilde{C}_{v_{i_0}, \dots, \hat{v}_{i_j}, \dots, v_{i_4}}$. This chain is in fact a 3-dimensional cycle in M^n , and it can be interpreted as a sphere of dim 2 in $Z_1(M^n, Z)$. This sphere is constructed as follows. Let $f_j : \bar{D}^2 \longrightarrow Z_1(M^n)$ be a map that corresponds to $(-1)^j \tilde{C}_{v_{i_0}, \dots, \hat{v}_{i_j}, \dots, v_{i_4}}$. Then, let $G_2 : \bar{D}^2 \longrightarrow Z_1(M^n)$ be a map that assigns to every $q \in \bar{D}^2$ a point $\sum_{j=0}^4 T_{f_j(x)}$. Then it is easy to see that the boundary of the disc is mapped to the zero 1-cycle, and we obtain a 2-sphere in $Z_{(60)}$. Now we want to use Proposition 4, so we need to lift this map of the 2-sphere to Γ_{60} . First we lift G_2 in the obvious way and consider $\tilde{G}_2 : \bar{D}^2 \longrightarrow \Gamma_{60}$. Next consider what happens to $\tilde{G}_2 : \partial\bar{D}^2 \longrightarrow \Gamma_{60}$. We see that each point is mapped to the union of 30 pairs of segments. Each pair consists of the same segment with different orientation. Those segments can be continuously contracted to their middles, (namely contract $\gamma \cup -\gamma$ to $\gamma(0.5)$). Thus we obtain a homotopy between the original circle and the circle that passes through constant parametrized 1-cycles only. This circle corresponds to 60 circles on a manifold, which we want to contract. Now unlike the proof of Theorem 1 above we cannot assume that M^n is simply connected. But we can contract these circles using the following simple construction that will similarly work for all dimensions: Consider the disc $\tilde{G}_2 : \bar{D}^2 \longrightarrow \Gamma_{60}$. For any $p \in \bar{D}^2$ $\tilde{G}_2(p) = \{\gamma_1^p, \dots, \gamma_{60}^p\}$. For each p consider $\{\gamma_1^p(0.5), \dots, \gamma_{60}^p(0.5)\}$. This determines a 2-dimensional disc in Γ_{60} that passes only through constant parametrized 1-cycles. This disc corresponds to 60 discs on the manifold. Now our circles can all be contracted over these discs, which establishes a homotopy between $\partial\bar{D}^2$ and a point in Γ_{60} . Now we can construct the required map of the 2-dimensional sphere into Γ_{60} providing the desired lifting to Γ_{60} exactly as this was done in the proof of Theorem 1. Also, note that in our present situation maps F_j defined in the text of Proposition 4 are 60 maps of S^2 to M^n defined as follows: For p in the southern hemisphere $F_j(p)$ is defined as $(\tilde{G}_2(p))_j(0.5)$. (Here we identify the southern hemisphere with D^2 .) For p in the northern hemisphere between the equator and a certain parallel $F_j(p)$ is

constant on every meridian. (This stage corresponds to contracting oppositely directed pairs of segments to the points in the middle corresponding to $t = 0.5$.) Finally, F_j maps the part of the northern hemisphere north of this parallel using $(\tilde{G}_2(p))_j(0.5)$ again. So, up to a homotopy F_j maps both hemispheres of S^2 in the same way. Hence F_j is contractible by the obvious homotopy. Therefore we can apply Proposition 4: Since our 2-sphere passes only through sufficiently short parametrized 1-cycles and because of our assumption, it can be contracted through sufficiently short parametrized 1-cycles. The 3-disc contracting this sphere corresponds to a 4-chain in M^n filling the 3-cycle we started from. (Here we have used the Almgren correspondence explained in the previous section). And so on.


It becomes obvious, that we can go on like that until we “extend” F to the $(n + 1)$ -skeleton of W thereby obtaining a $(n + 1)$ -singular chain in M^n filling modulo 2 the fundamental homology class of M^n which is clearly impossible. The resulting contradiction proves the theorem. QED.

Important Remark. As the reader noticed, we had not really used any notions or theorems of the geometric measure theory in order to prove Theorems 1 and 2. In fact, we could restrict ourselves to the consideration of rather elementary spaces Γ_k of parametrized 1-cycles instead of $Z_1(M^n, \mathbb{Z})$, and to only use Lemma 3 (but not Proposition 4), since all maps to the space of (non-parametrized) cycles that we constructed in the course of proving Theorems 1, 2 lift to appropriate spaces Γ_k (possibly after a reparametrization). Yet we preferred to make our exposition as above in order to illustrate geometric measure-theoretic origin of many ideas used in the present paper.


Acknowledgements. The paper was partially written during the visit of the authors to IHES in Summer of 2001. A part of this paper was written when both authors were working at the Courant Institute of Mathematical Sciences in 2001-2002. The authors would like to thank IHES and the Courant Institute for their kind hospitality. Both authors gratefully acknowledge the partial support of their research by NSERC research grants. Regina Rotman gratefully acknowledges the partial support of her research by the NSF Post-doctoral Fellowship.



(a)



(b)



(c)

REFERENCES

- [A] F.J. Almgren, Jr., *The homotopy groups of the integral cycle groups*, Topology **1** (1962), 257-299.
- [BM] E. Bierstone and P. Milman, *Semianalytic and subanalytic sets*, IHES Publ. Math. **67** (1988), 5-42.
- [B] M. Buchner, *Simplicial structure of real analytic cut loci*, Proc. Amer. Math. Soc. **64** (1977), 118-121.
- [BZ] Yu. D. Burago and V.A. Zalgaller, *Geometric inequalities*, Springer (1988).
- [C] C.B. Croke, *Area and the length of the shortest closed geodesic*, J. Diff. Geom. **27** (1988), 1-21.
- [ClCo] E. Calabi and J. Cao, *Simple closed geodesics on convex surfaces*, J. Diff Geometry **36** (1992), 517-549.
- [G] M. Gromov, *Filling Riemannian manifolds*, J. Diff Geometry **18** (1983), 1-147.
- [Ma] M. Maeda, *The length of a closed geodesic on a compact surface*, Kyushu J. Math. **48:1** (1994), 9-18.
- [Mi] J. Milnor, *Morse theory*, Princeton University Press (1963).
- [M] F. Morgan, *Geometric measure theory. A beginner's guide. Third Edition*, Academic Press (2000).
- [NR] A. Nabutovsky and R. Rotman, *The length of the shortest closed geodesic on a 2-dimensional sphere*, IMRN **2002:23** (2002), 1211-1222.
- [NR1] A. Nabutovsky and R. Rotman, *Upper bounds on the length of a shortest closed geodesic and quantitative Hurewicz theorem*, submitted for publication.
- [P] J. Pitts, *Existence and regularity of minimal surfaces on Riemannian manifolds*, Ann. Math. Studies **27** (1981), Princeton University Press.
- [R] R. Rotman, *Upper bounds on the length of the shortest closed geodesic on simply connected manifolds*, Math. Z. **233** (2000), 365-398.
- [S1] S. Sabourau, *Filling radius and short closed geodesics of the sphere*, preprint.
- [S2] S. Sabourau, *Global and local volume bounds and the shortest geodesic loop*, preprint.
- [SY] R. Schoen and S.-T. Yau, *Lectures on Differential Geometry*, International Press (1994).
- [Y] S.-T. Yau, *Problem section in "Seminar in Differential Geometry"*, ed. by S.-T. Yau, Princeton University Press (1982).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO, M5S 3G3, CANADA.

E-mail address: alex@math.toronto.edu and rina@math.toronto.edu